



Research Article

Evaluation of *Lablab purpureus* accessions in two agro-ecologies of Benishangul-gumuz, Western Ethiopia

Mulisa Faji^{1*}, Alemayehu Abebe², Kedija Ahmed², Diribi Mijena³, Habtamu Alebachewu²

¹Holetta Agricultural Research Center, Ethiopian Institute of Agricultural Research, Addis Ababa, Ethiopia

²Assosa Agricultural Research Center, Ethiopian Institute of Agricultural Research, Assosa, Ethiopia

³Wondogenet Agricultural Research Center, Ethiopian Institute of Agricultural Research, Shashemene, Ethiopia

Abstract

A study was conducted to evaluate *L. purpureus* accessions for their agronomic traits under two agro-ecologies of Benishangul-Gumuz region. The experiment was carried out at Assosa and Kamash forage research stations of Assosa Agricultural Research Center. They were purposively selected to represent mid and lowland agro-ecologies respectively. Four *Lablab purpureus* accessions (*L. purpureus* 147, 11609, 11640, and 6529) were evaluated in randomized complete block design with three replications. The general linear model procedures of SAS and least significance difference for data analysis and mean separation were employed respectively. Environment (E) had significant effect on plant height at forage harvesting ($P < 0.001$), forage dry matter yield ($P < 0.01$) and leaf to stem ratio ($P < 0.01$). *L. purpureus* genotypes planted at Kamash had longer plant height and dry matter yielder than Assosa, however leaf to stem ratio was higher at Assosa. Genotype (G) and G x E (G and E interaction) had no significant $P > 0.05$ effect on forage dry matter yield, plant height and leaf to stem ratio. In conclusion, environment had significant effect on forage dry matter yield, leaf to stem ratio and plant height, therefore evaluations of yield performance and adaptation patterns of *L. purpureus* genotype in multiple environments are very important step in agronomic evaluation and selection of better adapted and high yielding genotype.

Keywords: Accessions; Benishangul-Gumuz; Environment; *Lablab purpureus*

* Corresponding author: mulisa.faji2016@gmail.com

<https://doi.org/10.59122/134FA68>

Received May 10, 2022; Accepted October 10, 2022; Published December 20, 2022

© 2022 Arba Minch University. All rights reserved.

1. Introduction

Ethiopia has huge livestock population. However, it was not possible to meet the ever-increasing demand for animal products (Sisay, 2006). In Ethiopia, livestock production is characterized by low productivity for meat production and reproductive performance. The low productivity of livestock is mainly attributed to inadequate nutrition, low genetic potential of

indigenous breeds, prevalence of disease and parasites. Among these factors, under nutrition/malnutrition is considered to be the most important limiting factors constraining animal production in Ethiopia (Adugna et al., 2000; Bekele, 2010). In almost all parts of Ethiopia, annual pasture grasses (54.6%) and crop residues (31.6%) are the main sources of feed in the country (Tolera et al., 2000; Mengistu, 2004; CSA, 2017). Indeed, in Benishangul-Gumuz regional state natural grass land which is dominated by *Hypernia* species is the major livestock feed resource. Feedstuff of such composition, natural grass and crop residue, are insufficient to provide adequate quantity and quality of nutrient beyond maintenance requirement. Therefore, improved forage development and wise utilization of crop by-products can be minimizing the problem.

Cultivated forage crops have high yielder and better quality as compared to the natural pasture. Moreover, forage legumes have high protein sources that will enhance the activity of rumen microorganisms, increase intake and digestibility when supplement for animals consuming poor quality forages. Moreover, they can contribute to soil fertility maintenance and improvement (Mosi & Butterworth, 1985; Smith et al. 1989). Growing of multipurpose forage legume crops like *lablab* through under sowing/intercropping is useful for introduction of the forage crop and to use the available small farmland for both crop and livestock feed production. Intercropping annual forage legumes with cereal crops has been proposed as a strategy to control erosion, suppress weeds and contribute biological N to companion crop (Nadeem et al, 2009). Hence with this in mind, to introduce this forage species through intercropping forage development strategy, screening of the promising accessions to that environment is mandatory. The performance of forage species varies across locations due to differences in soil types, temperature and amount and distribution of rainfall. The cultivation of high-quality forages with a high yielding ability, adaptable to biotic and abiotic environmental stresses is one of the possible options to increase livestock production under smallholder farmers conditions (Tessema, 1999). Indeed, improved forage legume crops like *L. purpureus* are one of the best forage legumes for animal feed and soil fertility improvement through biological nitrogen fixation.

L. purpureus is an erect, short-lived perennial herbaceous crop often grown as an annual (Kikafunda, 2001). It is preferred for good forage production, nutritional quality, and palatability. *L. purpureus* is also shade-tolerant so that it is suitable for intercropping. *L. purpureus* is drought tolerant, and has been grown in arid and semiarid regions (Aganga &

Tshwenyane, 2003). Currently, *L. purpureus* is one of the major leguminous forage and green manure crops. However, there are no recommended *L. purpureus* accessions for their yield and agronomic performance in Benishangul-Gumuz region. Therefore, the objective of this study was to evaluate the forage dry matter yield production potential of *L. purpureus* accessions and finally select and recommend the best performing accessions for use as alternative improved forage legumes in Benishangul-Gumuz regional state (BGRS), Western Ethiopia and agro-ecologies similar to the study area.

2. Materials and Methods

2.1. Study areas

The trial was conducted under field conditions at Assosa, and Kamash forage research stations of Assosa Agricultural Research Center during the main cropping seasons. The test locations represent the mid and lowland areas ranging in altitude from 1000 to 1550 m.a.s.l. The farming system of the study areas is Agro pastoral. Descriptions of the test environments are indicated in Table 1.

Table 1. Descriptions of the test environments for geographical position

Parameters	Study sites	
	Assosa	Kamash
Latitude	10°30'N	09. °30'N
Longitude	034°20'E	35°45'E
Altitude(masl)	1500-1550	1000-1350
Annual rainfall(mm)	1316	1150
Daily minimum Temperature (°C)	16.75	25
Daily maximum Temperature (°C)	27.9	30

2.2. Experimental treatments and design

The four accessions of *Lablab purpureus* accessions (*L. Purpureus* 147, 11609, 11640, and 6529) were collected from ILRI. The accessions were planted in 3 m x 4 m plot using a randomized complete block design (RCBD) with three replications at the beginning of the main rainy season. Seed was sown by drilling in rows spaced 30 cm between rows, at a depth of 3 cm. Individual plot size was 12 m² and spacing between plots and replications 1.5 and 2 m, respectively at each testing environments. The treatments were sown according to their recommended seeding rates: 15-20 kg ha⁻¹.

2.3. Data collection

Data was collected on plant height at harvesting, leaf to stem ratio and forage dry matter yield. Plant height at harvesting was taken on six plants randomly selected from each plot and measured using a steel tape from the ground level to the highest leaf. For determination of biomass yield, accessions were harvested at 10% blooming stage using a quadrant which has 1m² areas. Weight of the total fresh biomass yield was recorded from each plot in the field and 500 g sample was taken from each plot to the laboratory and a sub-sample of known fresh weight was oven-dried for 72 hours at a temperature of 65 °C to determine dry matter yield.

2.4. Statistical analysis

Analysis of variance (ANOVA) procedures of SAS general linear model (GLM) was used to analyze the quantitative data (SAS, 2002). LSD test at 5% significance was used for comparison of means. The data was analyzed using the following model:

$$Y_{ijk} = \mu + G_i + E_j + GE_{ij} + B_k + e_{ijk}$$

Where, Y_{ijk} = dependent variables, μ = grand mean, G_i = effect of genotype i , E_j = effect of environment j , j = Assosa and Kamash, GE_{ij} = is the interaction effect of genotype i and environment j , B_k = effect of block k , and e_{ijk} = random error effect of genotype i , environment j , interaction effect of genotype i and environment j , and block k .

3. Results and Discussion

3.1. Environment and interaction effect on *L. purpureus* accessions performance

Combined analysis of variance for measured agronomic traits of *L. purpureus* accessions tested over environments is presented in Table 2. Environment (E) had significant effect on plant height at forage harvesting ($P < 0.001$), forage dry matter yield ($P < 0.01$) and leaf to stem ratio ($P < 0.01$). This variation might be due to differences among the testing environments in altitude, soil types, temperature and differences in both amount and distribution of annual rainfall and other agro-climatic factors. Findings from this experiment agree with reports of Abuye et al. (2018) who findings reveal that location significantly affect the forage dry matter yield of *L. purpureus* cultivars studied for herbage yield. Genotype (G) and G x E (genotype by environment interaction) had no significant ($P > 0.05$) effect on plant height at forage harvesting, leaf to stem ratio and forage dry matter yield. Generally, the result of this study indicated that environmental factors significantly influenced the yield performance and adaptation patterns of *L. purpureus* genotype. Therefore, evaluations of yield performance and adaptation patterns of *L.*

purpureus genotype in multiple environments are very important step in agronomic evaluation and selection of better adapted and high yielding species and varieties.

Table 2. Combined analysis of variance for measured agronomic traits of four *L. purpureus* accessions tested in two agro-ecologies of Benishangul-Gumuz regional state

Traits	Mean square		$G \times E$	Mean	Coefficient of variation (%)
	Genotype (G)	Environment (E)			
Plant height (cm)	ns	***	ns	145.21	44.09
Forage DM yield (t/ha)	ns	**	ns	4.34	36.89
Leaf to stem ratio	ns	**	ns	0.67	46.87

3.2. Leaf to stem ratio

The leaf to stem ratio at forage harvesting of tested *L. purpureus* accessions is indicated in Table 3. Environment had significant ($P < 0.01$) effect on leaf to stem ratio. The higher leaf to stem ratio was obtained from the mid-altitude (Assosa) than lowland altitude (Kamash) and this might be due the function of the longer periods of physiological growth of plants in high altitude/cooler environment with increase defoliation frequency stimulating leaf growth at the expense of stem production. The leaf to stem ratio has significant implications on the nutritive quality of the forage as leaves contain higher levels of nutrients and less fiber than stems. The result indicated that the leaf to stem ratio is an important factor affecting diet selection, quality and intake of forage (Smart et al., 2004).

Table 3. Mean leaf to stem ratio of four *L. purpureus* accessions tested in two agro-ecologies of Benishangul-Gumuz regional state

Accessions	Location/Environments		Combined analysis
	Assosa	Kamash	
<i>L. purpureus</i> 147	0.75	0.50	0.58
<i>L. purpureus</i> 11609	0.86	0.46	0.59
<i>L. purpureus</i> 6529	0.79	0.63	0.68
<i>L. purpureus</i> 11640	0.74	0.62	0.66
Mean	0.79 ^a	0.55 ^b	0.63
Coefficient of variation (%)	29.12	32.63	52.33
P-value	Ns	ns	ns

The leaf to stem ratio is associated with high nutritive value of the forage because leaf is generally of higher nutritive value (Tudsri et al., 2002) and the performance of animals is closely related to the amount of leaf in the diet. Therefore, according to the report of these authors the *L. purpureus* genotypes grown at Assosa is more nutritious than *L. purpureus* genotypes grown at Kamash due to the leaf to stem ratio was higher at Assosa than Kamash. The result of combined

and each locations analysis showed that leaf to stem ratio was non-significant ($P < 0.05$) among the accessions.

3.3. Plant height at forage harvesting

Summary analysis of variance for the performance of different *L. purpureus* accessions on plant height at forage harvesting was given in Table 4. The result of this study revealed that testing environment significantly ($P < 0.001$) affect plant height at forage harvesting and higher value of plant height at forage harvesting recorded at Kamash. This might be associated to the difference in environmental condition such as rainfall, soil fertility, temperature. Although, the result could be due to the Assosa soil is more acidic ($\text{PH} > 5.5$) than the Kamash soil ($\text{PH} < 5.5$) and this leads to plant root growth which result stunted and reduced growth rate in plants at Assosa. The result of combined and each location analysis showed that plant height at forage harvesting was not significantly ($P > 0.05$) different among the tested genotypes.

Table 4. Mean plant height at forage harvesting (cm) of four *L. purpureus* accessions tested in two agro-ecologies of Benishangul-Gumuz regional state

Accessions	Location/Environments		Combined analysis
	Assosa	Kamash	
<i>L. purpureus</i> 147	114.75	172.20	153.05
<i>L. purpureus</i> 11609	95.92	170.69	145.77
<i>L. purpureus</i> 6529	145.42	194.05	177.84
<i>L. purpureus</i> 11640	102.00	166.65	145.10
Mean	114.52 ^b	175.90 ^a	155.44
Coefficient of variation (%)	32.47	28.00	44.51
P-value	Ns	Ns	Ns

3.4. Forage and dry matter yield

The mean forage dry matter yield of four tested *L. purpureus* accessions is presented in Table 5. The result of this study showed that forage dry matter yield of *L. purpureus* was significantly ($P < 0.01$) different among locations/testing environment. *L. purpureus* genotypes grown at Kamash provided more yield than *L. purpureus* genotypes grown at Assosa. This might be due to the leaf stem ratio of the tested *L. purpureus* accessions was higher at Assosa than Kamash. As the leaf to stem ratio increased, the forage dry matter yield decreased, however the nutrient content increased and this might be due the leaf part of the forage is more nutritious than stem. The result might be due to the forage dry matter yield is directly related to plant height at forage harvesting stage, because when the plant height at forage harvesting increases forage dry

matter yield also increases. The result of combined and each location analysis showed that forage dry matter yield did not significantly ($P > 0.05$) affected by genotypes. Comparable value of herbage dry matter yield (4.54 t/ha) compared to this study result was reported by Bikila et al. (2017) who studied three accessions of *L. purpureus* (*L. purpureus* 11614, 147 and 11640) for their adaptation performance under the semi-arid environments of Borana zone.

Table 5. Mean forage DM yield (t/ha) of four *L. purpureus* accessions tested in two agro-ecologies of Benishangul-Gumuz regional state

Accessions	Location/Environments		Combined analysis
	Assosa	Kamash	
<i>L. purpureus</i> 147	3.40	4.38	4.06
<i>L. purpureus</i> 11609	3.27	4.98	4.41
<i>L. purpureus</i> 6529	4.36	5.59	5.18
<i>L. purpureus</i> 11640	3.90	4.81	4.51
Mean	3.73 ^b	4.94 ^a	4.54
Coefficient of variation (%)	27.68	36.57	36.76
P-value	Ns	ns	Ns

4. Conclusion

Based on the results, it can be concluded that the environment can affect plant height at forage harvesting, leaf to stem ratio and forage dry matter yield of *L. purpureus*. Lowland altitude could be most favorable agro-ecology than mid-altitude for *L. purpureus* since the plant height and forage dry matter yield is higher at lowland agro-ecology. Therefore, it can be concluded that evaluations of yield performance and adaptation patterns of *L. purpureus* genotype in multiple environments are very important step in agronomic evaluation and selection of better adapted and high yielding species and varieties.

Conflict of Interest

The authors declare that there is no conflict of interest.

Acknowledgement

The authors are grateful to Ethiopian Institute of Agricultural Research for financial support and Assosa Agricultural Research Center staffs specially Livestock Research Process for their honestly and heartily assistance.

References

Abuye, T., Temesgen, W., Diribsa, M., Keba, W., Fekede, G., & Kumsa, A. (2018). Herbage yield potential, crude protein yield and feeding value of selected *Lablab purpureus* cultivars

- grown under sub-humid climatic condition of Western Oromia, Ethiopia. *International Journal of Advanced Agricultural Research*, 6, 93–100.
- Adugna, T., Merkel, R. C., Goetsch, A. L., Sahlu, T., & Negesse, T. (2000). Nutritional constraints and future prospects for goat production in East Africa. In R. C. Merkel, G. Abebe, & A. L. Goetsch (Eds.), *The opportunities and challenges of enhancing goat production in East Africa* (pp. 43–57). Debub University.
- Aganga, A. A., & Tshwenyane, S. O. (2003). Lucerne, lablab and *Leucaena leucocephala* forage, production and utilization for livestock production. *Pakistan Journal of Nutrition*, 2(2), 46–53.
- Bekele, W. (2010). Effect of substitution of concentrate mix with *Sesbania sesban* on feed intake, digestibility, body weight change, and carcass parameters of Arsi-Bale sheep fed grass hay (Unpublished master's thesis). Haramaya University, Haramaya, Ethiopia.
- Bikila, N., Bedasa, E., & Jaldessa, D. (2017). Review of completed research activities of livestock research directorate. In *Proceedings of the regional workshop on review of completed research activities of Oromia Agricultural Research Institute* (pp. 1–247). Oromia Agricultural Research Institute.
- CSA. (2017). Agricultural sample survey. Volume II: Report on livestock and livestock characteristics (Private peasant holdings) (Statistical Bulletin 585). Central Statistical Agency, Federal Democratic Republic of Ethiopia.
- Kikafunda, J. (2001). Legume fallows for maize-based cropping systems in East Africa: Screening legumes for adaptability, biomass and nitrogen production. In *Proceedings of the 7th Eastern and Southern Africa Regional Maize Conference* (pp. 1–15). International Maize and Wheat Improvement Center.
- Mengistu, A. (2004). Pasture and forage resource profiles of Ethiopia. Ethiopia/FAO.
- Mosi, A. K., & Butterworth, M. H. (1985). The voluntary intake and digestibility of diets containing different proportions of tef (*Eragrostis tef*) straw and *Trifolium tembense* hay when fed to sheep. *Tropical Animal Production*, 10, 19–22.
- Nadeem, M. A., Iqbal, Z., Ayoub, M., Mubeen, K., & Ibrahim, M. (2009). Effect of nitrogen application on forage yield and quality of forage maize sown alone and in mixture with legumes. *Pakistan Journal of Life and Social Sciences*, 7, 161–167.
- Sisay, A. (2006). Qualitative and quantitative aspects of animal feed in different agro-ecological areas of North Gonder (Unpublished master's thesis). Alemaya University, Dire Dawa, Ethiopia.
- Smart, A. J., Schacht, W. H., Moser, L. E., & Volesky, J. D. (2004). Prediction of leaf/stem ratio using near-infrared reflectance spectroscopy (NIRS): A technical note. *Agronomy & Horticulture Faculty Publications*, 39.
- Smith, T., Manyuchi, B., & Mikayiri, S. (1989). Legume supplementation of maize stover. In B. H. Dzowela, A. N. Said, A. Wendem-Agenehu, & J. A. Kategile (Eds.), *Utilization of research results on forage and agricultural by-product material as animal feed resources in Africa* (pp. 302–320). International Livestock Centre for Africa.
- Tessema, Z. (1999). Napier grass adapts well in North Western Ethiopia. *Agri Topia*, 14(1).

- Tudsri, S., Jorgensen, S. T., Riddach, P., & Pookpakdi, A. (2002). Effect of cutting height and dry season date on yield and quality of five Napier grass cultivars in Thailand. *Tropical Grasslands*, 36, 248–252.