

## Research Article

## Enhancing environmental sustainability and income diversification through agroforestry practices in the Dollo Watershed, Kamba Zuria District, Southern Ethiopia

Yohannes Dikola Dito<sup>1\*</sup>, Aynalem Gochera Sade<sup>1</sup>, Ayele Chora Yota<sup>1</sup>

<sup>1</sup>Department of Physics, College of Natural and Computational Sciences, Dire-Dawa University, Dire-Dawa, Ethiopia

### Abstract

This study evaluated the contributions of agroforestry to community welfare and environmental health in targeted regions. Data were collected from 231 farm households across five villages using close-ended and open-ended questionnaires, complemented by focus group discussions with community leaders male and female heads of households, and farmers with various experience levels. A relative scoring method was employed to rank preferred woody species, with results indicating that 90% of participating farmers integrated agroforestry with monocropping, while only 9% practiced non-agroforestry agricultural activities. Among the 32 most preferred tree species identified for agroforestry, 79% were native and 21% were exotic. The findings revealed that 91% of respondents viewed agroforestry as their primary source of income while the 9% relied on non-agroforestry agricultural activities. Furthermore, 51% of households believed that agroforestry enhances biodiversity compared to monoculture, and 66.67% recognized increased crop output as its main advantage. However, as data collection was confined to a specific timeframe, seasonal variations in agroforestry systems and income generation were not fully captured. This underscores the necessity for innovative extension services and proactive engagement from governmental and non-governmental organizations to enhance the role of agroforestry in improving rural livelihoods and the environment.

**Keyword:** Carbon sequestration; Community; Environmental sustainability; Income; Soil improvement

\*Corresponding author: [dikola395@gmail.com](mailto:dikola395@gmail.com)

<https://doi.org/10.59122/2234abc>

Received October 18, 2024; Accepted November 8, 2024; Published December, 2024

© 2025 Arba Minch University. All rights reserved.

### 1. Introduction

Agriculture is well-thought as backbone and the single largest sector of Ethiopian economy in terms of employment opportunities for 85% of the total population, accounts for about 90% of exports materials and supplies over 90% of the raw materials for the agro industries (Zenebe et al., 2011) and total 33.88% of its GDP (Plecher, 2020). However, ensuring food security for the Ethiopians is becoming a big challenge, as a result of the rapid population growth, rapid urbanization, negative consequences of climate change, increasing demand for agriculture and forest products and

civil conflict (Yigezu, 2021). The Ethiopian population will be projected to be 171.8 million at 2050 by increasing at a rate of 2.5% annually (Bekele and Lakew, 2014). This will increase the demand for agricultural and livestock products (Hemathilake and Gunathilake, 2022). To mitigate these, adoption of the agroforestry practices is important. Agroforestry is a traditional land use system that may contribute to the solution of many present and future environmental problems (pantera et al., 2021). Agroforestry is a sustainable land management practice that includes the deliberate integration of a woody component with an agricultural production in the lower story (Damianidis et al., 2021).

In recent years, agroforestry has gained recognition as a crucial practice for promoting environmental sustainability and enhancing community livelihoods, particularly in regions such as Ethiopia. Empirical evidence confirms that agroforestry adoption supports to the farming income by generating an assured income for the local community (Rosati et al., 2020; Tesfay, 2024). Study carried out by Mulugeta and Mabrate (2017) Gedeo's indigenous agroforestry practices provided 40% of Ethiopia's premium grade coffee. Current studies of Bussa and Feleke, (2020) result revealed that increased income source and food security values of agroforestry practices for farmers had positive impression on their living standard.

The novelty of this special issue is the contribution for the understanding of how agroforestry practices will become a pathway for sustainable agricultural land management, environmental benefit, and as such help link science to practice. Agroforestry a solution to overcome the biodiversity challenges in the world, while fulfilling the European and Global Biodiversity Targets (Mosquera-Losada et al., 2020). The effort by Mantzanas et al. (2021) contributes to our knowledge on intercropping perennial trees with cereal crops species, is a very promising practice for Mediterranean areas with traditional olive agroforestry systems. Agroforestry could reverse the climate change impacts in many different ways and restore and promote soil health (Bateni et al., 2021). Healthy soil is arguably one of the most critical resources for the health of natural and agro ecosystems so that they can sustain food production as well as provision of ecosystem services.

Agroforestry provides numerous provisioning, regulating, cultural and supporting ecosystem services and environmental benefits while promoting eco-intensification based on a more efficient use of the land resources. Nevertheless, there is only little published recent information on the contribution of agroforestry to the environment in general and on climate change, carbon sequestration and forest fires in particular. In this special issue a number of articles are included that provide a multidimensional environmental benefits that agroforestry provides to the environment. The novelty of this special issue is the contribution for the understanding of how agroforestry practices will become a way for sustainable agricultural land management, and provide

multidimensional benefits that agroforestry provides to the community and environment in the study area. While agroforestry is increasingly recognized for its potential benefits in various regions, there is a lack of comprehensive study focusing specifically on the Dollo Watershed and its unique environmental and socio-economic context.

## 2. Materials and Methods

### 2.1. Study site

The study was conducted in Dollo watershed in Kamba zuria district Southern Ethiopia. The study watershed lies within 39° 37" E and 9° 41 " N. Situated 607 km south west of the capital city, Addis Ababa. The watershed forms parts of Gamo highlands of Ethiopia which is the part of Omo basin. The watershed is characterized by diverse topographic and elevation ranges from 1647 to 2180m.a.s.l. The annual average temperature of the area is 19.7 °C; annual average rainfall is 1470 mm. The farming system is a rain fed system.

### 2.2. Sample size determination and sampling technique

A household sample size was determined (Eq. 1) based on Yamane developed method (Yemane, 1967).

$$n = \frac{N}{1+N(e^2)} = 547/6.47 = 231 \quad (1)$$

Where, N is total household, n is number of sample size and 'e' is precision level for this case 5%. Systematic random sampling was used for selecting the participants from the total households as the total list of households was available.

### 2.3. Data collections

Data were collected from January 2022 to April 2023. Primary data for the study were collected using close-ended and open-ended questionnaire. The watershed had a total population of 547 household farmers of which 231 were selected systematically from household. Household socio-economic characteristics, farmers' species preferences, and agroforestry contribution to income diversification and environmental benefit were collected through questionnaire, focus group discussion and key informant interviews.

#### 2.3.1. Key informant interviews

The purpose of the key informant interviews was to collect their views on the role of agroforestry to household income diversification. The key informants include kebele administration leader, model farmers, elders and young farmers and agriculture development agents. From five villages 40 key informants were selected for interviews.

### 2.3.2. Focus group discussions

Two focus group discussions were conducted using randomly selected kebele administrations members. Separation of the gender groups provided equal opportunity for women and men to elicit and check perceptions and opinions on the role of agroforestry on household income diversification. All interviewees were informed about the purpose, subjects, and reasons of the research. Their participation was voluntary. The presence of kebele managers and local agricultural development agents facilitated the discussions and communication between researchers and respondents. Each focus group discussion lasted 1 hour.

### 2.3.3. Questionnaire

The data were collected using a standard close-ended and open-ended questionnaire administered through face-to-face interviews. This questionnaire was filled for the same rural respondents who adopted agroforestry activities.

### 2.4. Data analysis

The data collected from the survey were first checked for completeness. The quantitative data was analyzed through descriptive statistical analysis such as frequency, mean and percentage. The qualitative data were summarized by using narrative analyses. Simple majority/Relative score was used to calculate species preference in the study area.

## 3. Results and Discussion

### 3.1. Socio-economic characteristics of the household

During household surveys, data on age, family size, land holding size, and educational status were collected. The mean age of the respondents was between 25 and 35 years. Most respondents were males (84.5%), and the majority of the respondents (77.38%) were married. Most of the respondents (62%) had undergone formal education, with the majority (23.81%) having completed primary education, few (21.43%) had attended secondary education. Majority of the respondents (66.67%) had land size between 0 and 0.5 hector (Table 1).

### 3.2. Agroforestry practice

According to the survey on agroforestry practices, the majority (95.2%) of the population engaged in home gardens, followed by parkland (64.3%), woodlots (60.7%), and grazing systems (36.9%), respectively (Figure 1). These findings are consistent with Berihanu et al. (2020), who found that in northern Ethiopia, parkland agroforestry (90%) was practiced alongside woodlots (7%), home gardens (3%), and other land uses. The majority of the crops integrated with multipurpose trees, included *Zea mays* (maize) and *Eragrostis teff* (teff). Key informants explained that home garden

agroforestry practices are the most dominant in the study area since they are simple to manage. Similarly, key informants raised that parkland agroforestry is next highly adopted practice because of providing extra yield and services to human and animals.

Table 1. Socio-demographic characteristics of the households

Households data	Variables	Frequency	Percentage
Sex	Male	195	84.5
	Female	36	15.5
Family size	4	44	19.04
	5	50	21.43
	6	41	17.85
	7	36	15.47
	8	22	9.52
	9	25	10.71
	10	14	5.95
Age	Below 25	52	22.62
	Between 25-35	83	35.73
	Between 35-45	55	23.8
	Above 45	41	17.85
Marital status	Single	25	11.92
	Married	179	77.38
	Divorce	11	4.76
	Widowed	17	7.14
Land holding size (hector)	Below 0.5	154	66.67
	Between 0.5-0.1	52	22.62
	Between 1-1.5	17	7.14
	Above 1.5	8	3.57
Educational status	Not read and write	88	38.09
	Read and write	39	16.67
	1-8th grade	55	23.81
	8-12th grade	50	21.43

### 3.3. Species preference among farmers

A total of 32 different tree species were identified as suitable candidates for agroforestry, with 21% categorized as exotic species and 79% classified as native species. From highest to lowest, these were the top 10 that a significant percentage of farmers preferred: *Persea americana* (93%), *Coffee arabica* (87%), *Cordia africana* (83%), *Croton macrostachyus* (73%), *Combretum molle* (57%), *Casuarina equisetifolia* (45%), *Olea africana* (43%), *Terminalia brownii* (39%), *Eucalyptus globulus* (38%) and *Eucalyptus grandis* (38%) from highest to lowest (Figure 2). This study is consistent with studies of Alao and Shuaibu, (2013) and Adewusi, (2006), which showed that fruit trees both native and exotic are the most preferred. This suggests that food is considered the

most important resource, and the fire wood species are also the second critical resources for human existence.




Figure 1. The dominant agroforestry practices in the study area

Furthermore, the study's findings are consistent with those of Nkurikiye et al. (2024), who found that farmers had a favorable preference for planting trees that boost maize yields through agroforestry. In addition to contributing to crop production, *Calliandra calothyrsus*, *Gliricidia sepium*, and *Senna spectabilis* are studied for their ability to conserve natural resources, safeguard the environment, and produce exceptional fuel wood (Kuyah et al., 2020). These results also coincided with those of Tazebew and Asfaw (2018), who found that the choice of farmers to cultivate native multipurpose trees on their properties is a factor in the ecological and financial utility of the species when it comes to agroforestry practices centered on coffee.

The preference rating for woody species was calculated using a relative score. Respondents were asked to rank the top nine woody species from most preferred to list preferred among the species they plant and keep. The following woody species were preferred in order of significance: *Persea americana* (93%), *Coffee arabica* (87%), *Cordia africana* (83%), *Croton macrostachyus* (73%), *Combretum molle* (57%). Their preference was were ordered based the species ability to serve multiple purposes, such as providing food, generating income, providing firewood, being used as building materials, serving to shed from sunlight, providing fodder, and enhancing soil fertility (Table 2). Woody plants, both native and alien, are valuable assets on farms because of their significance to farmers' daily life. In the study area, *Cordia africana* and *Terminalia brownii* were the first and second species to integrate with crops. For example, species of trees that benefit agricultural crops, such as *Cordia africana* and *Terminalia brownii*, are planted widely throughout agriculture fields, but species of trees that compete with crops are planted individually to lessen their impact. Farmers set a variety of requirements for integrating trees on farmlands, such as the trees' ability to decompose quickly, their compatibility with crops, their multipurpose use value, their ability to promote soil fertility, and

their low branch volume. In the Lemo district of Southern Ethiopia, similar results were observed in crop livestock tree mixed systems (Kuria et al., 2014).




Figure 2. Species preference among farmers in the study area

The tree species found on grazing forms of agroforestry are incredibly large and dispersed, according to information from key informants. Based on information from key informants, field observations, and household interviews, the common tree species found in grazing land are *Terminalia schimperiana*, *Cordia Africana*, *Terminalia brownii*, *Ficus vasta*, *Croton macrostachyus*, and *Terminalia laxiflora*. According to data acquired from interviews with households, the types of trees recognized in agroforestry woodlots are large and densely populated. According to field observation and home interviews, the common tree species in woodlot agroforestry types are *Cupressus lusitanica*, *Eucalyptus grandis*, *Eucalyptus globulus*, *Combretum molle*, *Terminalia brownii*, and *Croton macrostachyus*. Agroforestry tree species that are found in woodlots are also listed.

Table 2. Respondents' species preference ranking according to their benefit in the study area

Species Scientific name	Respondents					Relative score					Total score	Rank	Reason of preference
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	5 <sup>th</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	5 <sup>th</sup>			
<i>Persea americana</i>	143	55	30	25	-	88.5	13.1	3.9	2.7	-	108.2	1 <sup>st</sup>	2, 1 and 5
<i>Coffee arabica</i>	55	55	25	74	14	13.1	13.1	2.7	23.7	0.85	53.65	2 <sup>nd</sup>	2,5 and 7
<i>Cordia Africana</i>	-	83	22	14	-	-	29.8	2.1	0.85	-	32.75	3 <sup>rd</sup>	2, 3 and 4
<i>Croton macrostachyus</i>	-	8	17	25	80	-	0.28	1.25	2.71	27.71	31.95	4 <sup>th</sup>	2, 3, 4 and 7
<i>combretum molle</i>	11	-	83	14	3	0.52	-	29.8	0.85	0.04	31.21	5 <sup>th</sup>	2, 4, 5 and 7
<i>causaria equistfolia</i>	-	8	28	22	28	-	0.28	3.39	2.09	3.39	9.15	6 <sup>th</sup>	3, 4, 6 and 7
<i>Olea Africana</i>	22	14	11	11	-	2.1	0.85	0.52	0.52	-	3.99	7 <sup>th</sup>	7, 2 and 1
<i>terminalia brownie</i>	-	-	-	22	3	-	-	-	2.09	0.04	2.13	8 <sup>th</sup>	5 and 7
<i>Eucalyptus globulus</i>	-	-	6	8	-	-	-	0.16	0.28	-	0.44	9 <sup>th</sup>	3 and 4
<i>Total</i>	231	223	222	215	128								

Footnote: relative score was calculated by multiplying the number of respondents in each ranks by its proportion (e.g. 143\*143/231=88.52). Reason of preference, 1= food, 2=income generation, 3= fire wood, 4= construction materials, 5= shade benefit, 6= animal fodder 7= soil fertility improvement (n=231)

The results of FGD, key informant interviews, and household surveys indicate that farmers' choices for particular species rely on the significance of those species within each type of agroforestry practice and how those components interact with one another. This conclusion is consistent with that of López-Sampson and Andrade (2024), who found that farmers placed a high value on animal temperature regulation and that providing environmental services can make agroforestry species more appealing. Furthermore, the findings of Hailu *et al.* (2024) investigation showed that farmers' choices for woody species differed across the nation and that they planted various woody species according to their respective advantages.

### 3.5. Role of agroforestry to livelihoods diversification

The farmers in the study area generate household income from both agroforestry and monoculture farms. Out of the farmers surveyed, 90% practice both agroforestry and monocropping. Meanwhile, 9% of the farmers engage in non-agroforestry farming activities, and the remaining 1% is involved in non-agricultural pursuits. Additionally, 91% of households reported that their primary source of income comes from agroforestry operations, while 9% indicated that their main source of income stems from non-agroforestry farming activities. These indicate that the agroforestry significantly increases farm income compared to non-agroforestry farm activities.



Table 3. Preferred tree species for different types of agroforestry in the Dollo watershed

Species scientific name	Home garden	Parkland	Grazing land	Woodlots	Percentage
<i>Annona muricata</i>	2	-	-	-	0.21645
<i>Carica papaya</i>	70	-	-	-	30.303
<i>Catha edulis</i>	12	-	-	-	5.19481
<i>Casuarina equisetifolia</i>	-	50	-	-	21.645
<i>Citrus sinensis</i>	13	-	-	-	5.62771
<i>Coffee Arabica</i>	110	-	-	-	47.619
<i>Combretum molle</i>	-	-	11	21	6.92641
<i>Cordia Africana</i>	0	0	36	23	6.38528
<i>Croton macrostachyus</i>	20	41	43	46	16.2338
<i>Cupressus lusitanica</i>	-	-	-	16	6.92641
<i>Ekebergia capensis</i>	-	12	-	-	5.19481
<i>Eucalyptus globulus</i>	-	-	-	34	14.7186
<i>Eucalyptus grandis</i>	-	-	-	31	13.4199
<i>Euphorbia ampliphyla</i>	-	-	-	19	8.22511
<i>Ficus vasta</i>	-	-	70	-	30.303
<i>Gossypium barbadense</i>	8	-	-	-	3.4632
<i>Grevillea robusta</i>	30	-	-	-	12.987
<i>Hypericum quartinianum</i>	-	-	-	14	6.06061
<i>Juniperus procera</i>	20	-	-	13	7.14286
<i>Mangifera indica</i>	60	23	-	-	17.9654
<i>Olea Africana</i>	-	32	7	-	8.44156
<i>Persea Americana</i>	120	98	-	-	47.1861
<i>Psydrax schimperiana</i>	-	-	-	18	7.79221
<i>Rhamnus prinoides</i>	60	-	-	-	25.974
<i>Schrebera alata</i>	-	-	9	-	3.8961
<i>Syzygium guineense</i>	-	-	-	19	8.22511
<i>Tamaranus indicus</i>	-	8	-	-	3.4632
<i>Terminalia brownie</i>	10	102	30	43	20.0216
<i>Terminalia laxiflora</i>	-	-	17	-	7.35931
<i>Terminalia schimperiana</i>	-	-	12	-	5.19481
<i>Vernonia amygdalina</i>	-	9	-	15	5.19481

Agroforestry can diversify Farmers' income in a number of ways. These include fruits, coffee, fodder crops, fire wood and dairy products. More than 93% of respondents use fruits from home garden agroforestry, such as banana, *Carica papaya*, *Persea americana*, and *Mangifera indica*; 90% use crops from parkland, such as *Zea mays* (Maize), *Eragrostis teff* (teff), haricot bean, and groundnuts; and 88% use woodlots, parkland, and home garden agroforestry for fuel and other wood products for construction. 83 % of the respondents declared that their primary

source of income comes from coffee plantations, which are a significant cash crop in the region. 67% of all respondents employed home gardens, parklands, and agroforestry to produce fodder for animals (Figure 3). Fruit was the most popular product harvested from agroforestry trees, demonstrating how dependent farmers were on these items for their primary source of revenue. This outcome was consistent with findings from related studies which looked into the roles of agroforestry in increasing farmers' income (Mabel *et al.*, 2017; Tharlakson, 2012; Quinon *et al.*, 2010). This outcome was consistent with reports from other academics (Kalaba, 2010; Maroyi, 2009) that said that agroforestry's various productions have enabled people to build sustainable lives.




Figure 3. List of products to livelihoods diversification from agroforestry practices

### 3.6. Role of agroforestry to environmental sustainability

Based on the results of the poll, farmers concur that agroforestry contributes to environmental sustainability by making more nutrients available to the soil, which helps to preserve and restore soil fertility. Out of all the households surveyed, 51% thought that agroforestry offered superior woody biodiversity than mono cropping farming, while 66.67% thought that the main advantage of agroforestry techniques was an increase in crop yield. Out of all the respondents, 46% claimed that agroforestry could improve soil nutrient availability and decrease soil erosion, while 36% claimed that it could provide shade for cattle and control the microclimate (Figure 4). Additionally, by increasing soil organic matter through leaf litter,

agroforestry practices improve soil fertility. Traditional land use practices like agroforestry could help find a solution of environmental problems in agriculture (Pantera *et al.*, 2021).




Figure 4. Contribution of agroforestry practices to environmental sustainability

One acknowledged effect of agroforestry practices was increased crop yield. This results from enhanced soil characteristics, microclimate, and nutrient levels (Fahad *et al.*, 2022). The current findings in lined with (Akinnesi *et al.*, 2006; Castle, 2021), who found that agroforestry techniques can raise crop yields in many regions of the world. Agroforestry in the Umbria area of Italy improved soil health and counteracted the negative impacts of soil erosion in a variety of ways (Bateni *et al.*, 2021; Pantera *et al.*, 2021). This result is consistent with that of Ndalama *et al.* (2015), who found that the primary ecosystem services received from agroforestry were soil improvement, water and nutrient retention and conservation, and biodiversity conservation. Increased agroforestry adoption, according to Khanal (2011), lessens the strain on forests and protected conservation areas. Furthermore, the present findings are consistent with the reviews conducted by Rolo *et al.* (2021) and Rosati *et al.* (2021), which suggested that implementing agroforestry techniques might enhance the sustainability of organic farming and augment soil

fertility. Additionally, Tsegaye's (2023) study result shows that agroforestry practices provide protective services such soil improvement, climate regulation, biological conservation, and recreational value in addition to their productive role.

#### 4. Conclusions

The study demonstrates that agroforestry significantly enhances the livelihoods and environmental sustainability of farmers. Practices such as home gardens, parklands, woodlots, and grazing systems are common, with a preference for multipurpose tree species like *Persea americana*, *Coffea arabica*, and *Cordia africana*. Agroforestry contributes to household income through products like fruits, coffee, and firewood, and improves soil quality, crop yields, and reduces soil erosion, aligning with previous research on its benefits for economic stability, biodiversity, and environmental resilience. To maximize agroforestry benefits, further research on species selection, management practices, and innovations is encouraged. Knowledge sharing platforms should be established to disseminate best practices. Integrating agroforestry into national and local agricultural policies can promote food security and income generation, especially in rural areas, through technical support, subsidies, and incentives. Agroforestry is vital for sustainable land management, enhancing soil fertility, and reducing soil erosion. It supports biodiversity conservation by promoting native, multipurpose species, contributing to ecosystem resilience. By providing diverse income sources, agroforestry reduces dependency on single crops and enhances resilience to market fluctuations and climate change, mitigating its impacts through improved soil health and carbon sequestration. Policymakers should consider agroforestry in climate adaptation and mitigation strategies, particularly in vulnerable regions.

#### Acknowledgment

The authors acknowledge Arba Minch University, College of Agriculture, for sponsoring this study.

#### Conflict of Interest

The authors declare no conflict of interest.

#### References

- Adewusi, S. R., Falade, M. S., Oyedapo, B. O., Rinaudo, T., & Harwood, C. (2006). Traditional and *Acacia colei* seed-incorporated diets in Maradi, Niger Republic. *Nutrition and Health*, 18(2), 161-177.
- Alao, J. S., & Shuaibu, R. B. (2013). Agroforestry practices and concepts in sustainable land use systems in Nigeria. *Journal of Horticulture and Forestry*, 5(10), 156-159.
- Asfaw Z, Agren, GI (2007). Farmers' local knowledge and topsoil properties of agroforestry practices in Sidama, Southern Ethiopia. *Agroforestry Systems*, 71 (1): 35-48.

- Bateni, C., Ventura, M., Tonon, G., & Pisanelli, A. (2021). Soil carbon stock in olive groves agroforestry systems under different management and soil characteristics. *Agroforestry Systems*, 95, 951-91.
- Bekele, A. , & Lakew, Y. (2014). Projecting Ethiopian demographics from 2012–2050 using the spectrum suite of models. *Ethiop Public Health Association*, 4, 1-6.
- Bussa, B., & Feleke, K. (2020). Contribution of parkland agroforestry practices to the rural community livelihood and its management in Southern Ethiopia. *Humanities and Social Sciences*, 8(4), 104-111.
- Castle, S.E., Miller, D.C., Ordonez, P.J., Baylis, K., & Hughes, K. (2021). The impacts of agroforestry interventions on agricultural productivity, ecosystem services, and human well-being in low-and middle-income countries: A systematic review. *Campbell Systematic Reviews*, 17(2), e1167.
- Damianidis, C., Santiago-Freijanes, J. J., den Herder, M., Burgess, P., Mosquera-Losada, M. R., Graves, A., Papadopoulos, A., Pisanelli, A., Camilli, F., Rois-Díaz, M., & Kay, S. (2021). Agroforestry as a sustainable land use option to reduce wildfires risk in European Mediterranean areas. *Agroforestry Systems*, 95, 919-929
- Fahad, S., Chavan, S. B., Chichaghare, A. R., Uthappa, A. R., Kumar, M., Kakade, V., Pradhan, A., Jinger, D., Rawale, G., Yadav, D. K., & Kumar, V. (2022). Agroforestry systems for soil health improvement and maintenance. *Sustainability*, 14(22), 14877.
- Hailu, F., Derero, A.' & Aticho, A. (2024). The impact of land uses on the diversity and farmers' preferences for woody species in the selected highlands of Ethiopia. *Agroforestry Systems*, 1-22.
- Hemathilake, D. M. K. S. & Gunathilake, D. M. C. C. (2022). Agricultural productivity and food supply to meet increased demands. In *future foods* (pp. 539-553). Academic Press.
- Khanal, S. (2011). Contribution of agroforestry in biodiversity conservation and rural needs fulfilment: A case study from Kaski District (Unpublished Doctoral Dissertation), Tribhuvan University, Nepal.
- Kuria, A., Lamond, G., Pagella, T., Gebrekirstos, A., Hadgu, K., & Sinclair, F.L. (2014). Local knowledge of farmers on opportunities and constraints to sustainable intensification of crop- livestock-trees mixed systems in Lemo Woreda, SNNPR Region, Ethiopian Highlands.
- Kuyah, S., Sileshi, G. W., Luedeling, E., Akinnifesi, F. K., Whitney, C. W., Bayala, J., Kuntashula, E., Dimobe, K., & Mafongoya, P. L. (2020). Potential of agroforestry to enhance livelihood security in Africa. *agroforestry for degraded landscapes: Recent Advances and Emerging Challenges*, 1, 135-167.
- López-Sampson, A., & Andrade, H. J. (2024). Agroforestry systems in Latin America. *Agroforestry Systems*, 98(5), 1075-1078.
- Mantzanas, K., Pantera, A., Koutsoulis, D., Papadopoulos, A., Kapsalis, D., Ispikoudis, S., Fotiadis, G., Sidiropoulou, A., & Papanastasis, V. P. (2021). Intercrop of olive trees with cereals and legumes in Chalkidiki, Northern Greece. *Agroforestry Systems*, 95, 895-905.
- Mosquera-Losada, M. R., Santiago-Freijanes, J. J., Rigueiro-Rodríguez, A., Rodríguez-Rigueiro, F. J., Martínez, D. A., Pantera, A., & Ferreiro-Domínguez, N. (2020). The importance of agroforestry systems in supporting biodiversity conservation and agricultural production: a European perspective. In *reconciling agricultural production with biodiversity conservation* (pp. 243-258). Burleigh Dodds Science Publishing.

- Mulugeta, G., & Mabrate, A. (2017). Production and ecological potentials of Gedeo's indigenous agro-forestry practices in Southern Ethiopia. *Journal of Resources Development and Management*, 30(1), 68-76.
- Ndalama, E., Kamanga-Thole, G. and Missanjo, E. (2015). Agroforestry contribution to the improvement of rural community livelihoods in Balaka, Malawi. *International Journal of Forestry and Horticulture*, 1(1), pp.5-11.
- NISR. (2023). The fifth Rwanda population and housing census: Main indicators report. The National Institute of Statistics of Rwanda (NISR). Kigali, Rwanda.
- Nkurikiye, J.B., Uwizeyimana, V., Van Ruymbeke, K., Vanermen, I., Verbist, B., Bizoza, A., & Vranken, L. (2024). Farmers' preferences for adopting agroforestry in the Eastern Province of Rwanda: A choice experiment. *Trees, Forests, and People*, 16, 100592.
- Pantera, A., Mosquera-Losada, M. R., Herzog, F., & Den Herder, M. (2021). Agroforestry and the environment. *Agroforestry Systems*, 95(5), 767-774.
- Plecher, H. (2020). Ethiopia: share of economic sectors in the gross domestic product (GDP) from 2009 to 2019.
- Rolo, V., Rocés-Díaz, J.V., Torralba, M., Kay, S., Fagerholm, N., Aviron, S., Burgess, P., Crous-Duran, J., Ferreira-Dominguez, N., Graves, A., & Hartel, T. (2021). Mixtures of forest and agroforestry alleviate trade-offs between ecosystem services in European rural landscapes. *Ecosystem Services*, 50, 101318.
- Rosati, A., Borek, R. and Canali, S. (2021). Agroforestry and organic agriculture. *Agroforestry Systems*, 95,805-821.
- Tazebew, E. and Asfaw, Z. (2018). Tree species diversity, preference and management in smallholder coffee farms of Western Wellega, Ethiopia. *Journal of Biology, Agriculture and Healthcare*, 8(24), 43-53.
- Tesfay, H.M., Oettel, J., Lapin, K., & Negash, M. (2024). Plant diversity and conservation role of three indigenous agroforestry systems of Southeastern Rift Valley landscapes, Ethiopia. *Diversity*, 16(1), 64.
- Tsegaye, N. T. (2023). The impact of agroforestry practice on forest conservation and community livelihood improvement: A case of Buno Bedele Zone of west Ethiopia'S Chora district. *Environmental and Sustainability Indicators*, 17, 100229.
- Yemana, T. (1967). *Statistics, an introductory analysis* (2<sup>nd</sup> ed.). New York, U.S.A.
- Yigezu Wendimu, G. (2021). The challenges and prospects of Ethiopian agriculture. *Cogent Food & Agriculture*, 7(1), 1923619.
- Zenebe, G, Jesper S, Alemu M, Atlaw A. (2011). Climate change and the Ethiopian economy. A computable general equilibrium analysis. *Environment for Development, Ethiopia*.